

Lamisil (Terbinafine)

Dermatology Associates of West Texas, LLP

For questions or an appointment, call (806)797-1202

Lamisil is an antifungal medication used to treat fungal infections of the skin and nails. It generally does not interact with most medications but if you are on blood thinners, anti-seizure medications or cholesterol lowering medications – then you will possibly need closer attention. Mild symptoms may occur such as stomach upset or change in taste and should resolve in time.

Adverse Reactions:

Any medication can cause any type of reaction. If you develop any unusual symptoms – notify the doctor.

Watch for the following:

Chronic or severe **stomach pain**

Chronic diarrhea

Skin rash or mouth ulcers or hives

Loss of appetite

Extreme tiredness

Fever, chills, sore throat or **signs of infection**

Special issues:

*Do not take Lamisil if you are pregnant

*Notify the doctor if you have **liver problems**

*You will be asked to **check blood-work 3 weeks** after starting treatment

NOTE: This is not a comprehensive information sheet. If you desire complete drug information, contact the pharmacist or refer to the “package insert”.

Lamisil (Terbinafine)

Dermatology Associates of West Texas, LLP

For questions or an appointment, call (806)797-1202

Lamisil is an antifungal medication used to treat fungal infections of the skin and nails. It generally does not interact with most medications but if you are on blood thinners, anti-seizure medications or cholesterol lowering medications – then you will possibly need closer attention. Mild symptoms may occur such as stomach upset or change in taste and should resolve in time.

Adverse Reactions:

Any medication can cause any type of reaction. If you develop any unusual symptoms – notify the doctor.

Watch for the following:

Chronic or severe **stomach pain**

Chronic diarrhea

Skin rash or mouth ulcers or hives

Loss of appetite

Extreme tiredness

Fever, chills, sore throat or **signs of infection**

Special issues:

*Do not take Lamisil if you are pregnant

*Notify the doctor if you have **liver problems**

*You will be asked to **check blood-work 3 weeks** after starting treatment

NOTE: This is not a comprehensive information sheet. If you desire complete drug information, contact the pharmacist or refer to the “package insert”.